

PRODUCT MONOGRAPH  
INCLUDING PATIENT MEDICATION INFORMATION

Pr **CABLIVI**®

Caplacizumab for injection  
Powder for solution (11 mg)  
Intravenous or subcutaneous  
Professed  
Antithrombotic Agent  
(B01AX07)

sanofi-aventis Canada Inc.  
1755 Steeles Avenue West  
Toronto, ON  
M2R 3T4

Date of Initial Authorization:  
February 28, 2020

Date of Revision:  
May 3, 2022

Distributed by Sanofi Genzyme,  
a division of sanofi-aventis Canada Inc.  
1755 Steeles Avenue West  
Toronto, ON  
M2R 3T4

Submission Control Number: 257723

s-a version 3.1 dated January 04, 2024

Cablivi® is a trademark of Ablynx N.V., Belgium. Used under licence.

## RECENT MAJOR LABEL CHANGES

7 WARNINGS AND PRECAUTIONS	09, 2021
----------------------------	----------

### TABLE OF CONTENTS

*Sections or subsections that are not applicable at the time of authorization are not listed.*

<b>RECENT MAJOR LABEL CHANGES</b> .....	<b>2</b>
<b>TABLE OF CONTENTS</b> .....	<b>2</b>
<b>PART I: HEALTH PROFESSIONAL INFORMATION</b> .....	<b>4</b>
<b>1 INDICATIONS</b> .....	<b>4</b>
1.1 Pediatrics.....	4
1.2 Geriatrics.....	4
<b>2 CONTRAINDICATIONS</b> .....	<b>4</b>
<b>4 DOSAGE AND ADMINISTRATION</b> .....	<b>4</b>
4.1 Dosing Considerations .....	4
4.2 Recommended Dose and Dosage Adjustment .....	4
4.3 Reconstitution .....	5
4.4 Administration.....	5
4.5 Missed Dose .....	5
<b>5 OVERDOSAGE</b> .....	<b>6</b>
<b>6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING</b> .....	<b>7</b>
<b>7 WARNINGS AND PRECAUTIONS</b> .....	<b>7</b>
7.1 Special Populations.....	9
7.1.1 Pregnant Women .....	9
7.1.2 Breast-feeding .....	9
7.1.3 Pediatrics.....	9
7.1.4 Geriatrics .....	9
<b>8 ADVERSE REACTIONS</b> .....	<b>9</b>
8.1 Adverse Reaction Overview.....	9
8.2 Clinical Trial Adverse Reactions .....	10
8.3 Less Common Clinical Trial Adverse Reactions .....	12

8.5	Post-Market Adverse Reactions .....	12
<b>9</b>	<b>DRUG INTERACTIONS .....</b>	<b>12</b>
9.4	Drug-Drug Interactions .....	12
9.5	Drug-Food Interactions.....	12
9.6	Drug-Herb Interactions.....	13
9.7	Drug-Laboratory Test Interactions.....	13
<b>10</b>	<b>CLINICAL PHARMACOLOGY .....</b>	<b>13</b>
10.1	Mechanism of Action.....	13
10.2	Pharmacodynamics .....	13
10.3	Pharmacokinetics .....	13
<b>11</b>	<b>STORAGE, STABILITY AND DISPOSAL .....</b>	<b>14</b>
<b>12</b>	<b>SPECIAL HANDLING INSTRUCTIONS .....</b>	<b>14</b>
<b>PART II: SCIENTIFIC INFORMATION .....</b>		<b>15</b>
<b>13</b>	<b>PHARMACEUTICAL INFORMATION .....</b>	<b>15</b>
<b>14</b>	<b>CLINICAL TRIALS .....</b>	<b>15</b>
14.1	Trial Design and Study Demographics .....	15
14.2	Study Results .....	17
<b>15</b>	<b>MICROBIOLOGY .....</b>	<b>18</b>
<b>16</b>	<b>NON-CLINICAL TOXICOLOGY .....</b>	<b>18</b>
<b>PATIENT MEDICATION INFORMATION.....</b>		<b>20</b>
<b>INSTRUCTION FOR USE .....</b>		<b>25</b>

## PART I: HEALTH PROFESSIONAL INFORMATION

### 1 INDICATIONS

Cablivi (caplacizumab) is indicated for the treatment of adults with acquired thrombotic thrombocytopenic purpura (aTTP) in combination with plasma exchange (PE) and immunosuppressive therapy.

#### 1.1 Pediatrics

Pediatrics (< 18 years of age): No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use.

#### 1.2 Geriatrics

Geriatrics ( $\geq$  65 years of age): Clinical studies of Cablivi did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects.

### 2 CONTRAINDICATIONS

Cablivi is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING.

### 4 DOSAGE AND ADMINISTRATION

#### 4.1 Dosing Considerations

Treatment with Cablivi is an adjunct to plasma exchange and immunosuppressive therapy.

#### 4.2 Recommended Dose and Dosage Adjustment

Cablivi should be administered upon initiation of plasma exchange therapy as recommended below:

- First day of treatment: 11 mg bolus **intravenous** injection at least 15 minutes prior to a plasma exchange followed by an 11 mg **subcutaneous** injection after completion of plasma exchange on that day.
- Subsequent days of treatment during plasma exchange: daily 11 mg **subcutaneous** injection following plasma exchange.
- Treatment after plasma exchange period: 11 mg subcutaneous injections once daily for 30 days following the last daily plasma exchange. If after initial treatment course, sign(s) of persistent underlying disease such as suppressed ADAMTS13 activity levels remain present, treatment may be extended for a maximum of 28 days.

Discontinue Cablivi if the patient experiences more than 2 recurrences of aTTP, while on Cablivi. Cases of relapse have been reported shortly after discontinuation of Cablivi treatment especially in case of

unresolved underlying auto-immune disease (see 7 WARNINGS AND PRECAUTIONS, Relapse of aTTP after stopping or interrupting use of Cablivi).

Health Canada has not authorized an indication for pediatric use (see 1 Indications, 1.1 Pediatrics)

### 4.3 Reconstitution

Cablivi should be prepared and reconstituted before administration. Detailed instructions on preparation and administration are provided in the INSTRUCTIONS FOR USE.

**Table 1:Reconstitution**

Vial Size	Volume of Diluent to be Added to Vial	Approximate Available Volume	Concentration per mL
12.5 mg	1 mL	1 mL	11 mg/mL solution

Cablivi is for single use only. Any unused medicinal product or waste material should be disposed of in accordance with local requirements

Unopened vial: see 11 STORAGE, STABILITY AND DISPOSAL. Do not use after the expiry date that is stated on the carton and label.

Reconstituted solution

- Use the CABLIVI solution immediately. If not, use CABLIVI within 4 hours after reconstitution when stored in the refrigerator at 2°C to 8°C (see 11 STORAGE, STABILITY AND DISPOSAL and INSTRUCTIONS FOR USE)

### 4.4 Administration

Withhold treatment for 7 days prior to elective surgery, invasive dental procedures, or other invasive interventions (7 WARNINGS AND PRECAUTIONS, Patients undergoing surgery).

The first dose should be administered by intravenous push injection by a healthcare provider. Subsequent doses should be administered by subcutaneous injections into the abdomen (INSTRUCTIONS FOR USE).

Do not inject into the area around the navel and do not use the same abdominal quadrant for consecutive injections.

Patients or caregivers may inject the medicinal product after proper training in the subcutaneous injection technique as instructed in INSTRUCTIONS FOR USE.

In the absence of compatibility studies, Cablivi must not be mixed with other medicinal products. For intravenous administration, if using an IV line, the line can be flushed with 9% Sodium Chloride Injection, or Glucose Injection 5% (w/v).

### 4.5 Missed Dose

Daily dosing and treatment continuity are critical. However, if a dose of Cablivi is missed during the plasma exchange period, it should be given as soon as possible. If a dose of Cablivi is missed after the

plasma exchange period, it can be administered within 12 hours of the scheduled time of administration. Beyond 12 hours the missed dose should be skipped and the next daily dose administered according to the usual dosing schedule.

## **5 OVERDOSAGE**

In case of overdose, based on the pharmacological action of caplacizumab, there is the potential for an increased risk of bleeding.

Close monitoring for signs and symptoms of bleeding is recommended. If needed, the use of von Willebrand factor (vWF) concentrate could be considered to correct hemostasis.

For management of a suspected drug overdose, contact your regional poison control centre.
---

## 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

To help ensure the traceability of biologic products, including biosimilars, health professionals should recognise the importance of recording both the brand name and the non-proprietary (active ingredient) name as well as other product-specific identifiers such as the Drug Identification Number (DIN) and the batch/lot number of the product supplied.

**Table 2 – Dosage Forms, Strengths, Composition and Packaging**

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Intravenous (IV) or subcutaneous injection	Powder for solution (11 mg)	Sucrose, citric acid anhydrous, trisodium citrate dihydrate, polysorbate 80.

Cablivi may be available in the following formats:

- Single vial Pack, containing:
  - 1 vial of Cablivi
  - 1 pre-filled syringe containing 1 mL of sterile water for injection (diluent for Cablivi)
  - 1 sterile vial adapter
  - 1 sterile needle
  - 2 alcohol swabs
- Multi-vial Pack, containing:
  - 7 vials of Cablivi
  - 7 pre-filled syringes containing 1 mL of sterile water for injection (diluent for Cablivi)
  - 7 sterile vial adapters
  - 7 sterile needles
  - 14 alcohol swabs

## 7 WARNINGS AND PRECAUTIONS

### General

**Relapse of aTTP after stopping or interrupting use of Cablivi:** Cases of relapse of aTTP have been reported in the clinical trials, especially within 30 days after discontinuation of caplacizumab (see 8 ADVERSE REACTIONS, 8.2 Clinical Trial Adverse Reactions). Most of the relapses occurred between 4 to 11 days after stopping Cablivi. In most of the patients who had relapse within 30 days after discontinuation of Cablivi treatment, ADAMTS13 activity levels were <10% at the end of the caplacizumab treatment, indicating that the underlying immunological disease was still active at the time caplacizumab was stopped. Therefore, patients, especially those with ADAMTS13 activity <10% at or near the time of discontinuation of Cablivi, should be closely monitored for platelet counts and signs of aTTP for early diagnosis of relapse after stopping or interrupting use of Cablivi.

### Driving and Operating Machinery

There have been no studies to investigate the effect of caplacizumab on driving performance or the ability to operate machinery. Bleeding episodes may result in fatigue that could impair the ability to operate or drive machinery.

## Hematologic

**Bleeding:** Cablivi increases the risk of bleeding (see 8 ADVERSE REACTIONS). Cases of major bleeding, including life-threatening and fatal bleeding have been reported in patients receiving caplacizumab, mainly in those using concomitant anti-platelet agents or anticoagulants. Caplacizumab should be used with caution in patients with underlying conditions that may predispose them to a higher risk of bleeding.

Interrupt use of Cablivi if clinically significant bleeding occurs and then closely monitor platelet counts and signs of relapse of aTTP (see 7 WARNINGS AND PRECAUTIONS, Relapse of aTTP after stopping or interrupting use of Cablivi). If needed, vWF concentrate may be administered to rapidly correct hemostasis. If Cablivi is restarted, monitor closely for signs of bleeding.

### Concomitant use of oral anticoagulants or high dose heparin

The risk of bleeding is increased with concomitant use of Cablivi with drugs affecting hemostasis and coagulation. Initiation or continuation of treatment with oral anticoagulants (e.g., vitamin K antagonists or direct oral anticoagulants (DOAC) such as thrombin inhibitors or factor Xa inhibitors) or high dose heparin requires a benefit/risk assessment and close clinical monitoring.

### Patients with coagulopathies

Due to a potential increased risk of bleeding, use of Cablivi in patients with underlying coagulopathies (e.g. hemophilia, other coagulation factor deficiencies) must be accompanied by close clinical monitoring.

### Patients undergoing surgery

If a patient is to undergo elective surgery, an invasive dental procedure or other invasive interventions, the patient must be advised to inform the physician or dentist that they are using Cablivi and it is recommended to withhold Cablivi treatment and closely monitor platelet counts and other signs of relapse of aTTP for at least 7 days before the planned intervention and continue the monitoring until Cablivi treatment is resumed (see 7 WARNINGS AND PRECAUTIONS, Relapse of aTTP after stopping or interrupting use of Cablivi). The patient must also notify the physician who supervises the treatment with Cablivi about the planned procedure. After the risk of surgical bleeding has resolved, and Cablivi is resumed, monitor closely for signs of bleeding.

If emergency surgery is needed, the use of von Willebrand Factor concentrate is recommended to correct hemostasis.

## Hepatic/Biliary/Pancreatic

**Severe hepatic impairment:** No formal studies with caplacizumab have been conducted in patients with severe hepatic impairment and no data regarding the use of caplacizumab in these populations are available. Use of Cablivi in patients with severe hepatic impairment requires a benefit/risk assessment and close clinical monitoring due to a potential increased risk of bleeding.

## Renal

**Renal Impairment:** Caplacizumab is mainly eliminated from the kidney (see 10 CLINICAL PHARMACOLOGY, 10.3 Pharmacokinetics). However, no data in patients with renal impairment are available as no formal studies of Cablivi in these patients have been performed.

## **Reproductive Health: Female and Male Potential**

- **Fertility**

The effects of Cablivi on fertility in humans are unknown. Dedicated animal studies assessing the effects of caplacizumab on male and female fertility have not been performed. In a 13-week repeat-dose toxicity study in cynomolgus monkeys, no impact of caplacizumab on male and female fertility parameters was observed.

### **7.1 Special Populations**

#### **7.1.1 Pregnant Women**

There are no available data on Cablivi use in pregnant women. No conclusions can be drawn regarding whether or not Cablivi is safe for use during pregnancy. Cablivi should be used during pregnancy only if the potential benefits to the mother outweigh the potential risks, including those to the fetus. All patients receiving Cablivi, including pregnant women, are at risk for bleeding. Monitor pregnant women and neonates for any evidence of excessive bleeding.

#### **7.1.2 Breast-feeding**

There are no available data on the presence of caplacizumab in human milk, effect on milk production, or the effects on the breastfed infant. No conclusions can be drawn regarding whether or not Cablivi is safe for use during breastfeeding. Cablivi should be used during breastfeeding only if the potential benefits to the mother outweigh the potential risks, including those to the breastfed child.

#### **7.1.3 Pediatrics**

The safety and efficacy of Cablivi in pediatric patients has not been established.

#### **7.1.4 Geriatrics**

Clinical studies of Cablivi did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients.

## **8 ADVERSE REACTIONS**

### **8.1 Adverse Reaction Overview**

The safety of Cablivi was evaluated in two placebo-controlled clinical studies (HERCULES [Phase III], in which 71 patients received Cablivi; and TITAN [Phase II], in which 35 patients received Cablivi). The data described below reflect exposure to Cablivi during the blinded periods of both studies, which include 106 patients with aTTP who received at least one dose, age 18 to 79 years, of whom 69% were female and 73% were White. The median treatment duration with Cablivi was 35 days (range 1-77 days).

The most frequently reported adverse reactions (>15%) were epistaxis, headache and gingival bleeding. Many of the reported adverse reactions were bleeding related. Serious bleeding adverse reactions were reported in  $\geq 2\%$  patients included epistaxis (4%) and subarachnoid hemorrhage (2%). Seven patients (7%) in the Cablivi group experienced an adverse reaction leading to study drug

discontinuation. None of the adverse reactions leading to discontinuation were observed in more than 1% of patients.

## **8.2 Clinical Trial Adverse Reactions**

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

Adverse reactions that occurred in  $\geq 2\%$  of patients treated with CABLIVI and more frequently than in those treated with placebo across the pooled data from the two clinical trials are summarized in **Table 3**.

**Table 3: Adverse Reactions in ≥ 2% of patients treated with caplacizumab and more frequent than placebo during the blinded periods of the Phase II and III aTTP studies (Safety Population) :**

<b>Adverse reaction</b>	<b>Cablivi (caplacizumab) (N=106) n (%)</b>	<b>Placebo (N=110) n (%)</b>
<b><i>Gastrointestinal disorders</i></b>		
Gingival bleeding	17 (16)	3 (3)
Rectal hemorrhage	4 (4)	0 (0)
Abdominal wall hematoma	3 (3)	1 (1)
<b><i>General disorders and administration site conditions</i></b>		
Fatigue	16 (15)	10 (9)
Pyrexia	14 (13)	12 (11)
Injection site hemorrhage	6 (6)	1 (1)
Injection site pruritus	3 (3)	0 (0)
<b><i>Musculoskeletal and connective tissue disorders</i></b>		
Myalgia	6 (6)	2 (2)
<b><i>Nervous system disorders</i></b>		
Headache	22 (21)	15 (14)
<b><i>Renal and urinary disorders</i></b>		
Hematuria	4 (4)	3 (3)
<b><i>Reproductive system and breast disorders</i></b>		
Vaginal hemorrhage	5 (5)	2 (2)
Menorrhagia	4 (4)	1 (1)
<b><i>Respiratory, thoracic and mediastinal disorders</i></b>		
Epistaxis	31 (29)	6 (6)
Dyspnea	10 (9)	5 (5)
<b><i>Skin and subcutaneous tissue disorders</i></b>		
Urticaria *	15 (14)	7 (6)

\*Urticaria: was observed primarily during the plasma exchange period

Ten patients in HERCULES experienced relapse during the 1-month follow up period (see 14 CLINICAL TRIALS). However, relapse only occurred in the Cablivi group (6/72, 8.3%) and in the patients who switched from placebo to open-label Cablivi due to exacerbation (3/25, 12%) but not in patients who only received placebo (0/48) (see 14 CLINICAL TRIALS). All these relapses except one occurred from 4 to 11 days after the last dose of Cablivi. One of these patients in the Cablivi group experienced severe thrombocytopenia and cerebral ischemia 6 days after stopping Cablivi treatment and died 2 days later; the death was considered not related to caplacizumab treatment. In 8 of these 9 patients, ADAMTS13 activity levels were <10% at the end of the study drug treatment, indicating that the underlying immunological disease was still active at the time Cablivi was stopped.

In TITAN, recurrences of TTP within 30 days of the last day of initial daily PE (i.e., exacerbations)

occurred in 8% of subjects (3/36) in the Cablivi group compared with 28% subjects (11/39) in the placebo group. During the first 30 days after stopping the study drug, relapse occurred in 22% (8/36) of the patients in the Cablivi group and none occurred in the placebo group (0/39). Most of these relapses (7/8) occurred within 10 days after discontinuation of Cablivi. In this study, treatment extensions beyond 30 days after stopping PE were not allowed and in most of the patients who had relapse, ADAMTS13 activity levels were <10% at the end of Cablivi treatment.

### ***Immunogenicity***

Immunogenicity assay results are highly dependent on several factors including assay sensitivity and specificity, assay methodology, sample handling, timing of sample collection, concomitant medications and underlying disease. For these reasons, comparison of incidence of antibodies to Cablivi with the incidence of antibodies to other products may be misleading.

Treatment emergent anti-drug antibodies (TE ADA) were detected in 3.1% of Cablivi treated patients in the HERCULES study. TE ADA were characterized as having neutralizing potential. There was no apparent impact on clinical efficacy or safety in the study.

### **8.3 Less Common Clinical Trial Adverse Reactions**

The following adverse drug reactions were observed in less than 2% of patients and at least in 1% of patients treated with Cablivi and more frequently than with placebo during the blinded periods of the Phase II and III aTTP studies:

- Hematochezia
- Hemoptysis

### **8.5 Post-Market Adverse Reactions**

The following adverse reactions have been identified during post approval use of Cablivi. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to caplacizumab exposure.

General disorders and administration site conditions:

*Frequency Not Known:* injection site erythema

Blood and lymphatic system disorders:

*Frequency Not Known:* major bleeding including life-threatening and fatal events

## **9 DRUG INTERACTIONS**

### **9.4 Drug-Drug Interactions**

No drug-drug interaction studies have been conducted with caplacizumab.

No interaction studies evaluating use of caplacizumab with oral anticoagulants (e.g. vitamin K antagonists, direct oral anticoagulants [DOAC] such as thrombin inhibitors or factor Xa inhibitors) or high dose heparin have been performed (See 7 WARNINGS AND PRECAUTIONS).

### **9.5 Drug-Food Interactions**

Interactions with food have not been established.

## 9.6 Drug-Herb Interactions

Interactions with herbal products have not been established.

## 9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

# 10 CLINICAL PHARMACOLOGY

## 10.1 Mechanism of Action

Caplacizumab is a humanized bivalent Nanobody (antibody fragment). It targets the A1-domain of von Willebrand factor and inhibits the interaction between von Willebrand factor and platelets, thereby reducing both vWF-mediated platelet activation and adhesion. As such, caplacizumab prevents the formation of ultra-large von Willebrand factor-rich platelet microthrombi but increases the risk of bleeding. It also affects the disposition of von Willebrand factor, leading to reduction in the level of both von Willebrand factor and factor VIII during treatment.

## 10.2 Pharmacodynamics

In the clinical trials in patients with aTTP (see 14 CLINICAL TRIALS), 10 mg subcutaneous dose of caplacizumab quickly elicited full inhibition of vWF-mediated platelet aggregation, as evidenced by RICO activity levels of < 20% approximately 4 hours post-dose and throughout the treatment period. RICO activity returned to baseline values within 7 days of drug discontinuation. In addition, 30-50% decrease in plasma vWF antigen levels and about 40% decrease in FVIII levels were also observed in these patients within one day after the first dose of caplacizumab and throughout the treatment course. The level of vWF antigen and FVIII returned to baseline value within 7 days of drug discontinuation.

The observed full inhibition of plasma vWF activity accompanied by large amount of decrease in serum vWF antigen and FVIII shortly after the first dose of caplacizumab and thereafter during the treatment period suggests that binding of caplacizumab to vWF accelerates the elimination of vWF and vWF-bound FVIII from the blood circulation. The caplacizumab-induced fast elimination of vWF and vWF-bound FVIII from blood circulation was dose-dependent as shown in clinical studies in healthy subjects.

## 10.3 Pharmacokinetics

Pharmacokinetics of caplacizumab appear as non-dose proportional, as characterized by target-mediated (vWF-mediated) disposition (see 10 CLINICAL PHARMACOLOGY, 10.2 Pharmacodynamics). Following a single subcutaneous dose of 10 mg caplacizumab to healthy subjects the mean (CV%) peak concentration (C<sub>max</sub>) was 528 (20%) ng/ml and AUC<sub>0-24</sub> was 7951(16%) hr·ng/ml. Following subcutaneous dosing of 10 mg caplacizumab daily for 14 days to healthy subjects, the mean (CV%) C<sub>max</sub> was 348 (30%) ng/ml and AUC<sub>0-τ</sub> was 6808 (26%) hr·ng/ml.

### **Absorption:**

After subcutaneous administration, caplacizumab is rapidly and almost completely absorbed (estimated F > 0.901) in the systemic circulation. In healthy volunteers receiving 10 mg caplacizumab subcutaneously once daily, the maximum concentration was observed at 6 - 7 hours post-dose and steady-state was reached following the first administration, with minimal accumulation in blood circulation.

**Distribution:**

After absorption, caplacizumab binds to vWF to form caplacizumab-vWF complexes. Excess free caplacizumab and caplacizumab-vWF complexes distribute to well perfused organs. In patients with aTTP the central volume of distribution was estimated at 6.33 L.

**Metabolism/Elimination:**

Due to small molecular weight (28 kD), free caplacizumab is assumed mainly cleared in the kidney. Caplacizumab-vWF complexes are heterogeneous (see 10 CLINICAL PHARMACOLOGY, 10.2 Pharmacodynamics). They are assumed to be proteolytically degraded especially by the reticuloendothelial system, including liver. The rate of the degradation may depend on the type of caplacizumab-vWF complexes and the tissue location. Higher levels of vWF, such as in patients with aTTP, increase the fraction of vWF-bound caplacizumab retained in the circulation. The t<sub>1/2</sub> of caplacizumab is, therefore, concentration and target (vWF) level-dependent.

**Antidrug Antibodies:**

No clinically significant differences in the pharmacokinetics of caplacizumab were observed in patients with pre-existing or treatment-emergent anti-drug antibodies.

**11 STORAGE, STABILITY AND DISPOSAL**

Store Cablivi in a refrigerator (2°C to 8°C).

Do not freeze.

Store in the original package in order to protect from light.

Do not use beyond the expiration date.

Cablivi may be stored at a temperature up to 30°C for a single period of up to 2 months, but not beyond the expiry date. Do not return Cablivi to refrigerated storage after storage at room temperature.

After the injection, throw away (dispose of) the used vial (with any remaining Cablivi liquid in it) with the adapter attached and the syringe with the needle attached in a sharps disposal container. See “Step 13: Throw away (dispose of) the used syringe and vial” at the end of the Instructions for Use for more disposal information.

**12 SPECIAL HANDLING INSTRUCTIONS**

Use the mixed Cablivi solution immediately. The mixed Cablivi solution can be stored for up to 4 hours in the refrigerator at 2°C to 8°C.

Please refer to the INSTRUCTIONS FOR USE.

## PART II: SCIENTIFIC INFORMATION

### 13 PHARMACEUTICAL INFORMATION

#### Drug Substance

Proper name: Caplacizumab

Chemical name: Caplacizumab

Molecular mass: 27876 Da

Structural formula:



Physicochemical properties: Purified Nanobody formulated in a citrate buffer (pH 6.5) containing sucrose polysorbate 80

#### Product Characteristics

Caplacizumab is a von Willebrand factor (vWF)-directed antibody fragment that consists of two identical humanized building blocks, linked by a three-alanine linker. Caplacizumab is produced in *Escherichia coli* by recombinant DNA technology and has an approximate molecular weight of 28 kDa.

### 14 CLINICAL TRIALS

#### 14.1 Trial Design and Study Demographics

The efficacy and safety of caplacizumab in addition to daily plasma exchange (PE) and immunosuppression in adults experiencing an episode of aTTP were established in a pivotal phase III trial study, "HERCULES" (Table 4).

**Table 4: Summary of patient demographics for clinical trials in acquired thrombotic thrombocytopenic purpura (aTTP)**

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n)	Mean age (Range)	Sex
ALX0681-C301  HERCULES	Phase III  Multi-centre; randomized; double-blind placebo-controlled	Caplacizumab or placebo: first day 11 mg iv bolus with PE followed by sc 11 mg injection. The 11 mg daily during daily PE period + 30 days. If needed treatment extension of daily 11 mg sc injections for 4x7 days.	145 patients	45 (18 – 79) years	100 females, 45 males

---

PE= plasma exchange  
i.v. = intravenous  
s.c. = subcutaneous

The efficacy of Cablivi (caplacizumab) in combination with plasma exchange and immunosuppressive therapy was established by HERCULES, a Phase III randomized double-blind placebo-controlled study in adult patients with an episode of aTTP (**Table 4**). A total of 145 patients were enrolled in the study; the median age was 45 (range: 18 to 79) years, 69% were female, 73% were White. After confirmation of eligibility to study participation and after the start of plasma exchange (PE) treatment, subjects were randomized in a ratio of 1:1 to receive study drug, i.e., Cablivi (n=72) or placebo (n=73) in addition to daily PE and immunosuppression. The maximum time allowed between the start of first PE (i.e., PE administered prior to randomization) and the start of the first PE after randomization (i.e., the first on-study PE) was 24 hours. Randomization was stratified according to the severity of neurological involvement (Glasgow Coma Scale score  $\leq 12$  or 13 to 15). Patients with sepsis, infection with *E. coli* 0157, atypical hemolytic uremic syndrome, disseminated intravascular coagulation or congenital thrombotic thrombocytopenic purpura were not eligible for enrollment:

The clinical trial protocol specified the Cablivi dose as 10 mg to be delivered by withdrawing all of the reconstituted solution from the vial and administering the full amount. A dose recovery study showed that the mean dose that can be withdrawn from a vial is 11 mg. Therefore, based on the dose recovery study, the mean dose delivered in the trial was 11 mg.

The Overall Study period in the study was composed of Study Drug Treatment period and a follow-up period. Recurrence of aTTP was defined as thrombocytopenia after initial recovery of platelet count (platelet count  $\geq 150 \times 10^9/L$  with subsequent stop of daily plasma exchange within 5 days), requiring re-initiation of daily PE. Recurrence of aTTP occurring during the first 30-days post-daily PE period, and recurrence of aTTP occurring after the 30-days post-daily PE period were termed as exacerbation and relapse, respectively.

In the Study Drug Treatment period, patients received a single intravenous bolus injection of 11 mg Cablivi or placebo from 6 hours to 15 minutes prior to the first PE on study. This was followed by daily subcutaneous injections of 11 mg Cablivi or placebo after completion of each plasma exchange for the duration of the daily PE period and for 30 days thereafter. Once the platelet count became  $\geq 150 \times 10^9/L$ , daily PE should continue for at least 2 days. If at the end of this treatment period there was evidence of persistent underlying immunological disease activity such as suppressed ADAMTS13 activity levels (indicative of an imminent risk for recurrence), treatment could be extended weekly for a maximum of 4 weeks together with optimization of immunosuppression.

Patients who experienced a recurrence while on study drug treatment were switched to open-label Cablivi. They were again treated for the duration of daily PE and for 30 days thereafter. If at the end of this treatment period there was evidence of ongoing underlying immunological disease, open-label treatment with Cablivi could be extended weekly for a maximum of 4 weeks together with optimization of immunosuppression.

In the follow-up (FU) period, patients were followed for 28 days after discontinuation of treatment. In case of recurrence during the follow-up period (i.e. after all study drug treatment had been stopped),

there was no re-initiation of Cablivi and the recurrence was to be treated according to the standard of care.

The median treatment duration with Cablivi was 35 days (range: 1 day to 77 days).

The primary endpoint was time to platelet count response, defined as initial platelet count  $\geq 150 \times 10^9/L$  with subsequent stop of daily PE within 5 days. It refers to the first time both conditions, platelet count above  $150 \times 10^9/L$  and the stop of daily PE within 5 days, were met.


A key secondary endpoint was the proportion of subjects with aTTP-related death, a recurrence of aTTP, or at least one treatment-emergent major thromboembolic event (e.g., myocardial infarction, cerebrovascular accident, pulmonary embolism or deep venous thrombosis [DVT]) during the Study Drug Treatment period.

Another key secondary endpoint was the proportion of subjects with a recurrence of aTTP in the Overall Study period.

## 14.2 Study Results

The Kaplan-Meier curves for time to platelet count response (the primary endpoint) are shown in **Figure 1**.

**Figure 1 - Platelet response over time**



Treatment with Cablivi resulted in a lower proportion of patients with aTTP-related death, recurrence of aTTP, or at least one treatment-emergent major thromboembolic event (a composite endpoint) during Study Drug Treatment Period (see **Table 5**).

**Table 5: Patients in HERCULES study with aTTP-related death, a recurrence of aTTP, or at least one treatment-emergent major thromboembolic event during Study Drug Treatment Period (ITT Population)**

	<b>Cablivi (caplacizumab) N=72</b>	<b>Placebo N=73</b>
<b>Number of patients with</b>	<b>n (%)*</b>	<b>n (%)</b>
TTP-related death	0	3 (4.1)
Recurrence of TTP (exacerbation)**	3 (4.2)	28 (38.4)
At least one treatment-emergent major thromboembolic event	6 (8.5)	6 (8.2)
<b>Total</b>	<b>9 (12.7)</b>	<b>36 (49.3)</b>
<b>p-value (CMH test adjusting for GCS at randomization)</b>	<b>&lt;0.0001</b>	

ITT = intent to treat population which includes all randomized patients  
 N = number of patients in the ITT population  
 n = number of patients with events  
 \* % based on 71 patients who received at least one dose of study drug  
 TTP = thrombotic thrombocytopenic purpura  
 \*\* recurrences during treatment period were considered exacerbations

The proportion of patients with a recurrence (an exacerbation or a relapse) of aTTP in the Overall Study period was 12.7% (9/71) in the Cablivi group compared to 38.4% (28/73) in the placebo group ( $p < 0.001$ ). In all 6 patients in the Cablivi group who experienced a recurrence of TTP during follow-up period (i.e. a relapse), ADAMTS13 activity levels were  $< 10\%$  at the end of the study drug treatment.

## 15 MICROBIOLOGY

No microbiological information is required for this drug product.

## 16 NON-CLINICAL TOXICOLOGY

### Repeat Dose Toxicology:

Toxicology studies have been conducted in guinea pigs and cynomolgus monkeys with doses yielding respective exposures of up to 50 and 24 times the expected exposure (AUC), respectively, from the daily human dose, and safety pharmacology was conducted as part of the repeat-dose studies.

Consistent with its mode of action, toxicology studies of caplacizumab have shown an increased bleeding tendency in guinea pigs (hemorrhagic subcutaneous tissue at the injection sites) and cynomolgus monkeys (hemorrhagic subcutaneous tissue at the injection sites, nose bleed, exaggerated menstrual bleeding, hematoma at sites of animal handling or experimental procedures, prolonged bleeding at injection sites). Furthermore, pharmacology-related decreases of von Willebrand factor

antigen, and consequently factor VIII:C, were noted in cynomolgus monkeys and, to a lesser extent for factor VIII:C, in guinea pigs.

In some monkeys anti FVIII antibodies (AFAs) against FVIII were observed following treatment with caplacizumab. In a 26-week repeated dose toxicity study, for two animals (one in the low and one in the mid dose groups), a transient anti-FVIII antibody response was noted on selected time points, which was absent during the recovery period. However, one male animal in the high dose group had marked decreases of the FVIII:Ag and FVIII:C during treatment that corresponded with a positive anti-FVIII antibody response resulting in bleeding leading to a scheduled sacrifice on day 85 of the study.

#### **Carcinogenicity and Mutagenicity:**

No studies have been performed to evaluate the mutagenic potential of caplacizumab, as such tests are not relevant for biologicals. Based on a carcinogenicity risk assessment, dedicated studies were not deemed necessary.

#### **Reproductive and Developmental Toxicology:**

##### Embryo-fetal development studies:

Two studies were conducted in pregnant guinea pigs with no reported signs of toxicity in dams or fetuses.

In an embryo-fetal development study, pregnant guinea pigs received caplacizumab via intramuscular administration in doses up to 20 mg/kg/day from the 6th to 41st day of pregnancy. No signs of systemic maternal toxicity were noted and no treatment-related influence was noted on body weight, body weight gain, net weight change from Day 6 onwards, or food and water consumption. No treatment-related influence was noted on the number of corpora lutea, implantation sites, resorptions, sex distribution, fetal and placental weights, number of live fetuses at birth, or the values calculated for the pre- and post-implantation loss when compared to the control.

A toxicokinetic study in pregnant guinea pigs assessed exposure of caplacizumab in the dams and fetuses. Caplacizumab was administered once daily to female guinea pigs at doses up to 40 mg/kg/day (corresponding to a drug exposure of 29 fold that observed in humans following a 11 mg/day dose) by intramuscular injection from the 6th to 41st day or 61st day of gestation. The results indicated exposure to caplacizumab in dams and, to a much lesser extent, fetuses, with no reported effects on fetal development. Fetal exposure to caplacizumab in primates and humans remains uncertain, as proteins lacking an Fc portion are not thought to freely pass the placental barrier.

##### Fertility:

Dedicated animal studies assessing the effects of caplacizumab on male and female fertility have not been performed. In repeat-dose toxicity studies in cynomolgus monkeys, no impact of caplacizumab on fertility parameters in male (testicular size, sperm function, histopathological analysis of testis and epididymis) and female (histopathological analysis of reproductive organs, periodic vaginal cytology) animals was observed.

## **PATIENT MEDICATION INFORMATION**

### **READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE**

Pr **CABLIVI**®

#### **Caplacizumab for injection, powder for solution**

Read this carefully before you start taking **Cablivi** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **Cablivi**.

#### **What is Cablivi used for?**

Cablivi contains the active substance caplacizumab. It is used to treat an episode of acquired thrombotic thrombocytopenic purpura (aTTP) in adults.

#### **How does Cablivi work?**

aTTP is a rare blood clotting disorder in which clots form in small blood vessels. These clots can block blood vessels and damage the brain, heart, kidneys, or other organs. Cablivi prevents the formation of these blood clots by stopping platelets in the blood from clumping together.

#### **What are the ingredients in Cablivi?**

Medicinal ingredients: caplacizumab

Non-medicinal ingredients: citric acid anhydrous, polysorbate 80, sucrose, trisodium citrate dihydrate

#### **Cablivi comes in the following dosage forms:**

Cablivi comes in a powder vial containing 11 mg of caplacizumab; boxes with 1 or 7 vials

#### **Do not use Cablivi if:**

You are allergic to caplacizumab or any of the other ingredients in this medication

#### **To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Cablivi. Talk about any health conditions or problems you may have, including if you:**

- bleed excessively or experience unusual symptoms such as headache, shortness of breath, tiredness, drop in blood pressure or fainting during treatment. Your doctor may ask you to stop the treatment. The doctor will let you know when you can start your treatment again
- are using anticoagulants (blood thinners) such as vitamin K antagonists, rivaroxaban, apixaban (which treat blood clots). Your doctor will decide how you should be treated
- are using anti-platelet medicines such as aspirin, or low molecular weight heparin (which prevent blood clots). Your doctor will decide how you should be treated

- have a bleeding disorder such as hemophilia. Your doctor will decide how you should be treated
- have severely reduced liver function. Your doctor will decide how you should be treated
- are going to have an operation, dental treatment or other procedures that required incisions. Your doctor will decide if it can be postponed or if you should stop Cablivi before your surgery or dental treatment
- are going to fully or temporarily stop taking Cablivi. There may be a risk of relapse of aTTP especially within 14 days after discontinuation of Cablivi treatment. Therefore, you should inform your doctor before you stop taking Cablivi and follow your doctor's instructions.

**Other warnings you should know about:**

Pregnancy and/or breastfeeding

Use of Cablivi is not recommended during pregnancy. If you are pregnant or planning to become pregnant, talk to your doctor about using Cablivi. Use of Cablivi is not recommended during breastfeeding. If you are breastfeeding or planning to breastfeed, talk to your doctor about using Cablivi.

Children and adolescents

Cablivi is not recommended for children under 18 years.

Cautions on driving and using machinery

Cablivi is not expected to influence the ability to drive or use machines. However, Cablivi-induced bleeding episodes may result in fatigue that could impair the ability to operate or drive machinery

**Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.**

**The following may interact with Cablivi:**

Also tell your doctor if you are using an anticoagulant (blood thinner) medicine such as vitamin K antagonists, rivaroxaban, or apixaban which treat blood clots or anti-platelet agents, such as aspirin, or low molecular weight heparin which prevent blood clots.

**How to take Cablivi:**

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

First Dose:

- 1 vial injected into a vein by a healthcare professional
- the medicine will be given before starting plasma exchange

Subsequent Doses (see INSTRUCTIONS FOR USE):

- 1 vial once daily as a subcutaneous injection (under the skin of the belly)
- the subcutaneous injection will be given after each daily plasma exchange
- after the daily plasma exchange finishes, your treatment with Cablivi will continue for at least

30 days with injection of 1 vial once daily.

- your doctor may ask you to continue daily treatment for up to additional 28 days
- follow your doctor's instruction after stopping Cablivi.

The first injection of Cablivi into your vein must be given by a healthcare professional. Your doctor may decide that you or your caregiver may inject Cablivi. In this case, your doctor or healthcare provider will train you or your caregiver on how to use Cablivi. Do not try to inject Cablivi until you have been taught how to do so by a healthcare professional. Never use the kit for another injection. Follow the details INSTRUCTIONS FOR USE that are included in each package.

#### **Overdose:**

If you think you, or a person you are caring for, have taken too much Cablivi, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

#### **Missed Dose:**

It is important that you take Cablivi every day as directed by your doctor. If you miss a dose of Cablivi during the plasma exchange period, it should be giving as soon as possible

If a dose of Cablivi is missed after the plasma exchange period, it can be taken within 12 hours of the scheduled dose time. Beyond 12 hours, the missed dose should be skipped, and the next daily dose taken according to the usual dosing schedule.

#### **What are possible side effects from using Cablivi?**

These are not all the possible side effects you may have when taking Cablivi. If you experience any side effects not listed here, tell your healthcare professional.

*Very common (may affect more than 1 in 10 people)*

- bleeding gums
- fever
- tiredness
- headache
- nosebleeds
- hives

*Common (may affect up to 1 in 10 people)*

- bruises on the abdomen
- rectal bleeding
- injection site reactions (for example, itching and bleeding)
- muscle pain
- blood in urine
- excessive bleeding during periods
- vaginal bleeding
- shortness of breath

Not known (*frequency cannot be estimated from the available data*)

- redness in the area where injection is given
- bleeding that may be serious or life-threatening

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
<b>VERY COMMON</b>			
Nosebleeds	√		
<b>COMMON</b>			
Excessive bleeding during periods	√		

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

<p><b>Reporting Side Effects</b></p> <p>You can report any suspected side effects associated with the use of health products to Health Canada by:</p> <ul style="list-style-type: none"><li>• Visiting the Web page on Adverse Reaction Reporting (<a href="https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html">https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html</a>) for information on how to report online, by mail or by fax; or</li><li>• Calling toll-free at 1-866-234-2345.</li></ul> <p><i>NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.</i></p>
---

**Storage:**

Cablivi should not be used after the expiry date which is stated on the label and carton.

Cablivi should be stored in a refrigerator (2°C to 8°C) in its original package to protect from light.

Cablivi may be stored at a temperature up to 30°C for a single period of up to 2 months, but not beyond the expiry date. Do not return Cablivi to refrigerated storage after storage at room temperature.

Use the mixed Cablivi solution immediately. The mixed Cablivi solution can be stored for up to 4 hours in the refrigerator at 2°C to 8°C.

Keep out of reach and sight of children.

**If you want more information about Cablivi:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website the sanofi-aventis Canada website [www.sanofi.com](http://www.sanofi.com), or by calling 1-800-265-7927.

This leaflet was prepared by sanofi-aventis Canada Inc.

Last Revised: May 3, 2022

## INSTRUCTIONS FOR USE

### Pr CABLIVI<sup>®</sup> Caplacizumab for injection, powder for solution

**Read the Instructions for Use before using Cablivi.**

Keep these instructions for future use. If you have any further questions, you should ask your health care professional or call 1-800-265-7927.




#### **Important Information:**



- Make sure the name Cablivi appears on the carton and vial label.
- For each injection, one Cablivi vial is needed. Only use the vial one time.
- Only use the supplies that are provided in the carton to prepare your prescribed dose.
- Do not use Cablivi after the expiration date on the carton.



Do not reuse any of the supplies. After your injection, throw away (dispose of) the used vial with any remaining Cablivi liquid in it. Throw away (dispose of) the used vial with the adapter attached and the syringe with the needle attached in a sharps disposal container. See “Step 13: Throw away (dispose of) the used syringe and vial” at the end of this INSTRUCTIONS FOR USE for more disposal information.




#### **How do I Store Cablivi?**



- Store Cablivi in the refrigerator between 2°C to 8°C.
- Used the mixed Cablivi solution immediately. The mixed Cablivi solution can be stored for up to 4 hours in the refrigerator at 2°C to 8°C.
- Cablivi may be stored at a temperature up to 30°C for a single period of up to 2 months, but not beyond the expiry date. Do not return Cablivi to the refrigerator after it has been stored at room temperature.
- Do not freeze.
- Store in original package to protect from light.
- Keep out of reach and sight of children.



INSTRUCTIONS	RELATED IMAGE				
<p>Each Cablivi carton contains either of the following:</p>					
<table border="1"> <thead> <tr> <th data-bbox="175 432 451 485">Single vial Pack</th> <th data-bbox="451 432 740 485">Multi-vial Pack</th> </tr> </thead> <tbody> <tr> <td data-bbox="175 485 451 947"> <ul style="list-style-type: none"> <li>1 vial of Cablivi</li> <li>1 prefilled syringe containing 1 mL Sterile Water for Injection (diluent for Cablivi)</li> <li>1 sterile vial adapter</li> <li>1 sterile needle</li> <li>2 alcohol swabs</li> </ul> </td> <td data-bbox="451 485 740 947"> <ul style="list-style-type: none"> <li>7 vials of Cablivi</li> <li>7 prefilled syringes containing 1 mL Sterile Water for Injection (diluent for Cablivi)</li> <li>7 sterile vial adapters</li> <li>7 sterile needles</li> <li>14 alcohol swabs</li> </ul> </td> </tr> </tbody> </table>	Single vial Pack	Multi-vial Pack	<ul style="list-style-type: none"> <li>1 vial of Cablivi</li> <li>1 prefilled syringe containing 1 mL Sterile Water for Injection (diluent for Cablivi)</li> <li>1 sterile vial adapter</li> <li>1 sterile needle</li> <li>2 alcohol swabs</li> </ul>	<ul style="list-style-type: none"> <li>7 vials of Cablivi</li> <li>7 prefilled syringes containing 1 mL Sterile Water for Injection (diluent for Cablivi)</li> <li>7 sterile vial adapters</li> <li>7 sterile needles</li> <li>14 alcohol swabs</li> </ul>	<p style="text-align: center;"><b>Contents of CABLIVI carton</b></p>  <p style="text-align: center;"><b>Additional supplies needed (not in CABLIVI carton)</b></p> 
Single vial Pack	Multi-vial Pack				
<ul style="list-style-type: none"> <li>1 vial of Cablivi</li> <li>1 prefilled syringe containing 1 mL Sterile Water for Injection (diluent for Cablivi)</li> <li>1 sterile vial adapter</li> <li>1 sterile needle</li> <li>2 alcohol swabs</li> </ul>	<ul style="list-style-type: none"> <li>7 vials of Cablivi</li> <li>7 prefilled syringes containing 1 mL Sterile Water for Injection (diluent for Cablivi)</li> <li>7 sterile vial adapters</li> <li>7 sterile needles</li> <li>14 alcohol swabs</li> </ul>				
<ul style="list-style-type: none"> <li>Additional supplies needed: <ul style="list-style-type: none"> <li>Sharps disposal container (see “step 13: throw away (dispose of) the used syringe” at the end of this Instructions for Use for more disposal information.</li> <li>Cotton balls</li> </ul> </li> </ul>					
<p><b>Before preparing a dose of Cablivi</b></p> <ul style="list-style-type: none"> <li>Wash your hands well with soap and water</li> <li>Prepare a clean flat surface</li> <li>Check to make sure the carton contains all of the items needed to prepare a dose</li> <li>Check the expiration date (see Figure A). <b>Do not use Cablivi if the date has passed</b></li> <li><b>Do not use Cablivi if the packaging or any supplies inside of the carton are damaged in anyway</b></li> </ul>	 <p style="text-align: center;"><b>Figure A</b></p>				




INSTRUCTIONS	RELATED IMAGE
<p><b>Step 1: Bring the vial and syringe to room temperature</b></p> <ul style="list-style-type: none"> <li>Place all the supplies in the carton on the clean flat surface</li> <li>If the carton was not stored at room temperature, allow the vial and the syringe to reach room temperature by holding them in your hands for 10 seconds (See Figure B). <b>Do not use any other way to warm up the vial and syringe.</b></li> </ul>	 <p style="text-align: center;"><b>Figure B</b></p>
<p><b>Step 2: Clean the rubber stopper</b></p> <ul style="list-style-type: none"> <li>Remove the green plastic cap from the metal cap of the vial (See Figure C). <b>Do not use the vial if the green plastic cap is missing.</b></li> </ul>	 <p style="text-align: center;"><b>Figure C</b></p>
<ul style="list-style-type: none"> <li>Clean the exposed rubber stopper using one alcohol swab to wipe it and allow it to dry for a few seconds (See Figure D).</li> <li><b>After cleaning the rubber stopper, do not touch it or allow it to touch any surface.</b></li> </ul>	 <p style="text-align: center;"><b>Figure D</b></p>



INSTRUCTIONS	RELATED IMAGE
<p><b>Step 3: Attach the vial adapter</b></p> <ul style="list-style-type: none"> <li>• Take the vial adapter and remove the paper cover (See Figure E). Leave the vial adapter in its opened plastic packaging for now. <b>Do not touch the adapter itself.</b></li> <li>• <b>Place the adapter over the vial, while keeping the adapter in its plastic packaging.</b></li> </ul>	 <p style="text-align: center;"><b>Figure E</b></p>
<ul style="list-style-type: none"> <li>• Press down firmly on the adapter until it snaps into place, with the adapter spike pushing through the vial stopper (See Figure F). <b>Do not remove the adapter from the vial once attached.</b></li> <li>• <b>Keep the adapter in its plastic packaging.</b></li> </ul>	 <p style="text-align: center;"><b>Figure F</b></p>
<p><b>Step 4: Prepare the syringe</b></p> <ul style="list-style-type: none"> <li>• Pick up the syringe.</li> <li>• While holding the syringe with one hand, break off the white plastic cap by snapping at the perforation of the cap with your other hand (See Figure G).</li> <li>• <b>Do not use the syringe if the white plastic cap is missing, loose, or damaged.</b></li> <li>• <b>Do not touch the syringe tip or allow it to come into contact with any surfaces.</b></li> <li>• Lay the syringe on the clean flat surface.</li> </ul>	 <p style="text-align: center;"><b>Figure G</b></p>



INSTRUCTIONS	RELATED IMAGE
<p><b>Step 5: Connect the syringe to adapter and vial</b></p> <ul style="list-style-type: none"> <li>Remove the plastic packaging from the adapter attached to the vial by holding the vial with one hand, pressing the sides of the adapter packaging with your other hand, and then lifting the packaging upwards (See Figure H).</li> <li>Be sure that the adapter does not detach from the vial.</li> </ul>	 <p style="text-align: center;"><b>Figure H</b></p>
<ul style="list-style-type: none"> <li>Hold the adapter with the attached vial with one hand. Place the tip of the syringe on the connector part of the vial adapter using the other hand.</li> <li>Gently lock the syringe into the vial adapter by turning it clockwise <b>until it cannot twist any further</b> (See Figure I).</li> </ul>	 <p style="text-align: center;"><b>Figure I</b></p>
<p><b>Step 6: Prepare the solution</b></p> <ul style="list-style-type: none"> <li>Place the vial upright on the flat surface with the syringe pointing downwards.</li> <li>Slowly push the syringe plunger down until the syringe is empty (See Figure J). <b>Do not remove the syringe from the vial adapter.</b></li> </ul>	 <p style="text-align: center;"><b>Figure J</b></p>


INSTRUCTIONS	RELATED IMAGE
<ul style="list-style-type: none"> <li>With the syringe still connected to the vial adapter, gently swirl the vial, with syringe attached, until the powder is dissolved in the vial (See Figure K). <b>Do not shake the vial.</b></li> </ul>	 <p style="text-align: center;"><b>Figure K</b></p>
<ul style="list-style-type: none"> <li>Allow the vial with the attached syringe to stand on the flat surface for <b>2 minutes</b> at room temperature to allow the powder to completely dissolve (See Figure L). The plunger may rise up by itself again, this is normal.</li> </ul>	 <p style="text-align: center;"><b>Figure L</b></p>
<p><b>Step 7: Draw up solution</b></p> <ul style="list-style-type: none"> <li>Check the solution in the vial for particles, cloudiness, or clumps. All powder must be fully dissolved and the solution must be clear. <b>Do not use the medicine if you see particles, cloudiness, or clumps.</b> Use a new carton of Cablivi and call your healthcare provider if this happens.</li> <li>Slowly press the syringe plunger fully down.</li> <li>Keep the syringe on the vial and turn the whole vial, adapter and syringe- upside down.</li> <li>Slowly pull the plunger down to withdraw all of the solution from the vial into the syringe (See Figure M). <b>Do not shake it.</b></li> </ul>	 <p style="text-align: center;"><b>Figure M</b></p>

INSTRUCTIONS	RELATED IMAGE
<p><b>Step 8: Detach the syringe</b></p> <ul style="list-style-type: none"> <li>After drawing up the solution into the syringe, turn the whole - vial, adapter and syringe- right-side up (with the syringe at the top) and place on the flat surface (See Figure N).</li> </ul>	 <p style="text-align: center;"><b>Figure N</b></p>
<ul style="list-style-type: none"> <li>Detach the filled syringe from the adapter by holding the vial and adapter in one hand and gently twisting the syringe counter-clockwise with the other hand (See Figure O).</li> <li>Throw away the vial and the attached adapter into a sharps disposal container.</li> <li><b>Do not touch the syringe tip or allow it to touch the clean flat surface.</b> Place the syringe on the clean flat surface.</li> </ul>	 <p style="text-align: center;"><b>Figure O</b></p>
<p><b>Step 9: Attach the needle</b></p> <ul style="list-style-type: none"> <li>Open the needle package by using both thumbs to pull apart the packaging (See Figure P).</li> <li>Remove the needle with the needle cap from the package.</li> </ul>	 <p style="text-align: center;"><b>Figure P</b></p>

INSTRUCTIONS	RELATED IMAGE
<ul style="list-style-type: none"> <li>Attach the needle with the needle cap to the syringe by turning clockwise <b>until it cannot twist any further</b> (See Figure Q). <b>Do not remove the needle cap.</b></li> </ul>	 <p data-bbox="1040 640 1136 667">Figure Q</p>
<ul style="list-style-type: none"> <li>Pull back the needle safety shield (See Figure R).</li> </ul>	 <p data-bbox="1040 1085 1136 1113">Figure R</p>
<p><b>Step 10: Prepare your injection site</b></p> <ul style="list-style-type: none"> <li>Select an injection site on your stomach (abdomen) (See Figure S). Avoid the 2-inch area around your belly button (navel). Select a different injection site from the one you used on the previous day to help the skin to recover after the injection.</li> </ul>	 <p data-bbox="1040 1470 1136 1497">Figure S</p>

INSTRUCTIONS	RELATED IMAGE
<ul style="list-style-type: none"> <li>Clean your injection site with an alcohol swab (See Figure T). Let your skin dry.</li> </ul>	 <p data-bbox="1040 625 1133 653">Figure T</p>
<p data-bbox="188 688 516 716"><b>Step 11: Giving your injection</b></p> <ul style="list-style-type: none"> <li>Carefully remove the needle cap from the needle and throw it away in a sharps disposal container (See Figure U). Make sure the needle does not touch anything before the injection.</li> <li>Hold the syringe at eye level with the needle pointing upwards.</li> </ul>	 <p data-bbox="1040 1047 1133 1075">Figure U</p>
<ul style="list-style-type: none"> <li>Check to see if there are any air bubbles. If there are any air bubbles, remove them by tapping the side of the syringe with your finger until they rise towards the tip (See Figure V).</li> </ul>	 <p data-bbox="1040 1472 1133 1499">Figure V</p>

INSTRUCTIONS	RELATED IMAGE
<ul style="list-style-type: none"> <li>Then, slowly push the plunger up until a small amount of liquid drips from the needle (See Figure W).</li> </ul>	 <p data-bbox="1036 653 1138 680">Figure W</p>
<ul style="list-style-type: none"> <li>Gently use one hand to pinch the skin that has been cleaned between your thumb and forefinger, making a fold (See Figure X).</li> <li>Hold the pinch during the entire injection.</li> </ul>	 <p data-bbox="1040 1087 1136 1115">Figure X</p>
<ul style="list-style-type: none"> <li>Use your other hand to insert the needle all the way into your skin fold at a 45 to 90-degree angle (See Figure Y).</li> <li>Push down on the plunger of the syringe until all of the solution is injected into your skin.</li> <li><b>Pull out the needle at the same angle you inserted it. Do not rub your injection site.</b></li> </ul>	 <p data-bbox="1040 1577 1136 1604">Figure Y</p>

INSTRUCTIONS	RELATED IMAGE
<p><b>Step 12: After your injection</b></p> <ul style="list-style-type: none"> <li>• Right after your injection, move the needle safety shield over the needle until it clicks into place (See Figure Z).</li> <li>• In case you are bleeding at the injection site, place a cotton ball over the skin right away. Press gently on the cotton ball until the bleeding has stopped. If bleeding does not stop, call your healthcare provider.</li> </ul>	 <p style="text-align: center;"><b>Figure Z</b></p>
<p><b>Step 13: Throw away (dispose of) the used syringe and vial.</b></p> <ul style="list-style-type: none"> <li>• Throw away the syringe with the needle and the vial with the adapter in a sharps disposal container right away after use. Do not throw away (dispose of) loose needles and syringes with your household trash.</li> <li>• If you do not have a sharps disposal container, you may use a household container that is: <ul style="list-style-type: none"> <li>• made of a heavy-duty plastic,</li> <li>• can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out,</li> <li>• upright and stable during use,</li> <li>• leak-resistant, and</li> <li>• properly labeled to warn of hazardous waste inside the container.</li> </ul> </li> <li>• When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of your sharps disposal container.</li> <li>• Do not dispose of your used sharps disposal container in your household trash unless your community guidelines permit this. Do not recycle your used sharps disposal container.</li> </ul>	